



HAL
open science

Performance of wheat-based cropping systems and economic risk of low relative productivity assessment in a sub-dry Mediterranean environment

A. Nasrallah, Hatem Belhouchette, N. Baghdadi, M. Mhawej, T. Darwish, S. Darwich, G. Faour

► To cite this version:

A. Nasrallah, Hatem Belhouchette, N. Baghdadi, M. Mhawej, T. Darwish, et al.. Performance of wheat-based cropping systems and economic risk of low relative productivity assessment in a sub-dry Mediterranean environment. *European Journal of Agronomy*, 2020, 113, pp.125968. 10.1016/j.eja.2019.125968 . hal-02467539

HAL Id: hal-02467539

<https://hal-ciheam.iamm.fr/hal-02467539>

Submitted on 26 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 Performance of wheat-based cropping systems and economic
2 risk of low relative productivity assessment in a sub-dry
3 Mediterranean environment

4 A. Nasrallah^{a,b,c,*}, H. Belhouchette^a, N. Baghdadi^b, M. Mhaweji^c, T. Darwish^c,
5 S. Darwich^d, G. Faour^c

6 ^aCIHEAM-IAMM, UMR-System, 34090 Montpellier, France

7 ^bIRSTEA, University of Montpellier, UMR TETIS, 34090 Montpellier, France

8 ^cNational Center for Remote Sensing, National Council for Scientific Research (CNRS), Riad al Soloh, Beirut 1107 2260, Lebanon

9 ^dFaculty of Agriculture, Lebanese University, Beirut 99, Lebanon

10
11 **Abstract**

12 The promotion of optimum rotations and agricultural management of winter wheat-based
13 cropping systems is very critical, as wheat is considered an essential component in the
14 Mediterranean diet. Considering the delicate economic situation of farmers in the
15 Mediterranean area, recommending a low risk, sustainable farming system is desirable. In
16 this study, an innovative application of a multi-criteria field-level approach is presented,
17 targeting food security, farmer profitability and environmental sustainability. The
18 CropSyst biophysical simulation model was calibrated and implemented for the study site.
19 It was chosen for its agro-environmental robustness to simulate four rotations (wheat-
20 wheat, wheat-fallow, wheat-potato, and wheat-fava bean). Four types of wheat agricultural
21 management systems (full fertilization and full irrigation, full fertilization and zero
22 irrigation, zero fertilization and full irrigation, and zero fertilization and irrigation) were
23 tested in low and high soil water holding capacity (WHC) types. The effects of soil
24 conditions, management practices and rotation type on wheat grain yields were assessed.
25 Furthermore, the performance of each winter wheat-based cropping system was evaluated
26 in terms of productivity (protein production and profitability) and the efficient use of

27 resources (nitrogen and water), as well as the economic risk of low relative productivity
28 each one engenders. The results show that there is no particular optimal scenario that can
29 simultaneously ensure high productivity, reduce economic risk of low relative productivity,
30 and achieve high wheat- water- and nitrogen-use efficiency. However, the wheat-fava bean
31 rotation cultivated with no wheat fertilization appeared to be a better substitute to the
32 wheat-wheat rotation in terms of protein production (0.93 t/ha versus 0.8 t/ha in low WHC
33 soil and 1.34 t/ha versus 1.17 t/ha in high WHC). This cropping system achieved a higher
34 net profit (2111 US\$/ha versus 1222US\$/ha in low WHC and 3550 US\$/ha versus 2450
35 US\$/ha in high WHC), showing high resource-use efficiency and was less risky for
36 farmers. Moreover, a very high profit could only be attained with the wheat-potato rotation
37 (8640 US\$/ha and 12170 US\$/ha in low and high WHC, respectively), yet with low input-
38 efficiency and high economic risk of low relative productivity.

39 *Keywords:* Winter wheat, CropSyst, Risk, Efficiency, Management, Cropping system,
40 Lebanon.

41

42 **1. Introduction**

43 Throughout history, the Mediterranean, especially its eastern and southern parts, has been
44 known to be the origin of many landraces and a pioneer in food production. It has never
45 been a region of abundance and glut, yet has always overcome the deficiencies in
46 production (Braudel, 1990; Kehoe, 1988). Winter wheat (*Triticum Durum L.*) is one of the
47 major crops grown in the Mediterranean. In Lebanon and the Middle East and North Africa
48 (MENA) region, wheat is often financially and sometimes technically supported as a part
49 of the governmental subsidy system. This self-sufficiency policy has long been the bedrock

50 of food security, leading to the continuous cultivation and successive sowing of wheat (El
51 Khansa, 2017; Nasrallah et al., 2018). At the same time, the MENA region is the largest
52 cereal importer in the world, with over 58 million metric tons, covering more than 50% of
53 its consumption (Wright and Cafiero, 2010). For nations within the MENA region,
54 importing cereal grains (mainly wheat) is not a matter of choice, but a necessity (Ahmed
55 et al., 2014).

56 Even if policies and policy-makers are keen to encourage large cereal production (wheat
57 in particular), simple wheat-based cropping systems co-exist in the Mediterranean region
58 (MoA, 2010). Intensive local wheat production in monoculture (wheat-wheat rotations) has
59 always been coupled with drawbacks and nutrient mining and deficiency. According to
60 Sieling et al. (2005), wheat-following-wheat rotations indeed lead to reduced yields,
61 compared to wheat following a different crop. The main reasons behind this finding are (1)
62 the increase in biotic yield-limiting factors (Bennett et al., 2012) and (2) a lesser availability
63 of needed nutrients and particularly nitrogen (Dalal et al., 2001; Sieling et al., 2005).
64 Thereby, the different already existing wheat-based cropping systems (with different
65 rotation and management practices) are directly linked to soil water and nitrogen access
66 (Pala et al., 2007; Ryan et al., 2007), production type (e.g. cereal grains, legume grains or
67 vegetables), in addition to economic risk, which farmers can overcome (Komarek et al.,
68 2015; Sadras, 2002).

69 Thus, several studies have tried to address the obvious question of the performance and
70 outputs of each production system. Diverse crop rotations have been experimented, and
71 sometimes versus monoculture systems (Beaudoin et al., 2005; Constantin et al., 2010;
72 Hansen et al., 2015, 2010; Macdonald et al., 2005; Moreau et al., 2012; Sieling and Kage,

73 2006). For instance, long-term field experiments in Central and Western Europe have
74 shown that the inclusion of a catch crop within a rotation can indeed significantly increase
75 nitrogen-use efficiency (NUE) as well as the N (Nitrogen) uptake of the main crop
76 (Berntsen et al., 2006; Constantin et al., 2011). In comparing different types of wheat-based
77 rotations, Angus et al. (2015) found that both fallow-wheat and break crop-wheat rotations
78 generally produced greater yields than wheat-wheat rotations. For instance, legume-wheat
79 rotations generated over 20% wheat grain yields compared to wheat-wheat rotations.
80 Without underestimating the role of plant genetics, the efficient management of water and
81 N has been identified as a crucial need for closing the yield gap, which is estimated by
82 comparing the observed yield with the attainable one (Mueller et al., 2012), of main cereal
83 crops (Sinclair and Rufty, 2012) notably on arid and semi-arid soils with low organic
84 carbon and nitrogen content (Darwish et al., 2018). Downscaling to field and farm levels
85 made it possible to study and analyse the economic risk that farmers and producers could
86 face, in relation to their adopted cropping systems (Di Falco and Perrings, 2005; Komarek
87 et al., 2015; Mahmood et al., 2017; Valle et al., 2004). However, the absence of a clear
88 integrated approach at field level, assessing different existing wheat-based cropping
89 systems regarding their productivity, resource-use efficiency and economic risk of low
90 productivity, represent the motivation of this study. It raises the key issue of the wheat-
91 based cropping systems to be promoted (regarding resources, soil types, climatic
92 variability, and management systems). It also offers a conceptual guide-map, allowing
93 policy-makers and producers to categorize different cropping systems with reference to
94 productivity (i.e. net profit and protein production), efficiency (water and nitrogen) and the
95 economic risk of low relative productivity.

96 For this purpose, the biophysical simulation model "CropSyst" version 3 (Monzon et al.,
97 2012) was calibrated and evaluated in the mid-Bekaa plain in Lebanon based on extensive
98 field work. Scenarios concerning different existing wheat-based cropping systems (rotation
99 type and wheat management system) in two soil types with contrasting water holding
100 capacities were developed and run against historical long-period climatic data (i.e. 20
101 years). Based on the model outputs, the objectives of the paper were to first, to measure
102 and compare the effect of the different rotations (wheat-wheat, wheat-potato, wheat-fallow
103 and wheat-fava beans) and agricultural practices (water and nitrogen) on winter wheat
104 grain yield. Second, to evaluate and compare the performance of each cropping system (of
105 rotation type and agricultural practices) in terms of productivity and efficiency of utilizing
106 the resources. Finally, to establish the link between the performance of each cropping
107 system and its economic risk of low relative productivity.

108 **2. Methods**

109 **2.1. Study site and crop management**

110 The Bekaa plain in Lebanon is located between 33°33' N and 33°60' N latitude, 35°39' E
111 and 36°14' E longitude (Fig. 1). The area of the plain is around 860.25 km² with an average
112 elevation of 1000 m above sea level. The dominant soils within the plain are mainly clay
113 to loam but differ in their water holding capacity (WHC). The Bekaa is characterized by a
114 semi-arid Mediterranean climate and the average annual precipitation is around 600 mm.
115 In addition, agriculture is the main economic activity as field crops, orchards, annual and
116 perennial plants are cultivated.

117

118 Field crop areas (e.g. cereals, vegetables, alfalfa and legumes) range from 0.1 ha to 20 ha.
119 65% of the national cereal production is produced in the Bekaa plain, while winter wheat
120 areas in the plain correspond to 44% of the national wheat area, occupying areas ranging
121 from 9000 to 12000 ha annually. 51% of potato crops, which is one of the largest tuberous
122 crops cultivated in Lebanon, is cultivated in the Bekaa plain as one of the most important
123 cash crops. As for legumes, Bekaa is responsible for 20% of the national cultivation area,
124 16% of this area corresponds to fava beans, which occupy around 1548 ha in the plain
125 (MoA, 2010). Wheat and fava beans are winter crops, as they are sown in November, while
126 potatoes are sown in March.

127 Regarding irrigation management, 72% of Bekaa crops are fully or supplementary
128 irrigated. Even though fava beans and wheat are grown during the winter season, they
129 receive supplementary irrigation during early spring to ensure better yields, although 20%
130 of the wheat in the Bekaa plain is not supplementary irrigated (due to no access to water,
131 money shortage and/or in the hope of a good rainy season) (El Khansa, 2017). While
132 potatoes, on the other hand, are fully irrigated (on a weekly basis) from sowing to
133 harvesting, ranging from 10 to 20 mm per application, depending on the phenological stage
134 (Darwish et al., 2003, 2006a).

135 Fertilization is supplied, especially nitrogen, being one of the most growth driving nutrient.
136 Fertilization management practices differ among farmers, however, nitrogen is supplied in
137 both organic and inorganic forms. In the case of wheat, farmers supply nitrogen in amounts
138 of up to 230 kg ha⁻¹ as ammonium sulfate at emergence and before the flowering stages.
139 However, up to 15% of wheat farmers do not apply synthetic fertilization. This is mainly

140 due to the cultivation of potato as a previous crop, where the land is supposed to be fertile
141 enough to meet wheat nitrogen demands, besides other economic considerations and
142 money saving purposes, when necessary. As for potato crops, nitrogen is applied before
143 planting in the form of manure (around 250 kgN ha⁻¹), in addition to a second application
144 of nitrogen (around 100 kgN ha⁻¹) before the inception of flowering. When fertilized, fava
145 beans receive a triggering amount of 50 kgN ha⁻¹ of nitrogen 60 days after sowing.

146 In the Bekaa plain, one of the most followed rotation types is the wheat-potato rotation as
147 it is one of the most profitable rotations. In fact, the existence of wheat-legume rotations is
148 limited within the Bekaa plain. However, some farmers do cultivate wheat in monoculture
149 to benefit from governmental support in buying their harvest with relatively good prices.
150 23% of wheat cultivated land in 2016 had also been wheat cultivated in 2017 (Nasrallah et
151 al., 2018). Therefore, among suitable agriculture land, less than 1500 ha of land are left as
152 fallow annually, corresponding to 4% of the total exploited area.

153

154 **2.2.Simulation model**

155 **2.2.1. CropSyst model description**

156 The CropSyst model, which was first presented by Stockle et al. (2003, 1994), uses
157 weather, soil and crop input data for the estimation of crop productivity under different
158 management conditions (water and nutrient input). It has been widely applied to many
159 regions (e.g. USA, China, Central and Northern Europe and the Mediterranean) and crops
160 (e.g. cereals, vegetables and legumes) (Ahmad et al., 2017; Belhouchette et al., 2008;
161 Brooks et al., 2017; Palosuo et al., 2011; Rötter et al., 2012), especially for its ability to

162 work on a daily basis for simulating multi-crop scenarios, in addition to water-soil
163 dynamics (Richard's equation for our case) and nitrogen budgets.

164 **2.2.2. Datasets for model calibration and evaluation**

165 *Experimental datasets for winter wheat model calibration and evaluation*

166 Five winter wheat plots were selected within the region of mid-Bekaa, corresponding to
167 two dominant soil types with different water holding capacities (low: 100-175 mm/m and
168 high: 175-250 mm/m). Above ground biomass (AGB), dry matter production (DMP), soil
169 water content (SWC), and above ground nitrogen (AGN) were measured at four
170 physiological stages and replicated three times within each winter wheat plot (Table 1).

171

172

173 Above ground biomass (AGB) was measured by a destructive method. After weighing the
174 fresh samples for each replication within each plot, the samples were cut up, mixed and
175 quartered and a representative sample was oven dried at 70 °C until constant weight was
176 reached (Catchpole and Wheeler, 1992).

177 Soil water content (SWC) was measured using the gravimetric method. For each of the
178 three pedological horizons (the depth of each soil horizon varied among plots), a sample
179 of soil was taken out, weight measured fresh, then sent to the oven to dry at 105 °C until
180 constant weight was obtained. For each winter wheat reference plot, the measurement was
181 replicated three times randomly at each depth at five crop development stages (Reynolds,
182 1970).

183 Above ground nitrogen (AGN) was measured following the Kjeldahl N method (Rodriguez
184 and Miller, 2000). Crop N uptake for each treatment was calculated based on the
185 corresponding data of dry matter production and N content for each treatment, at each
186 sampling date. Winter wheat in-situ measurements are summarized in table 2. It shows the
187 minimum, maximum and average results of the measurements of replications. SWC
188 measurements in the different horizons were also considered.

189

190

191 *Survey datasets for potato and fava bean model calibration and evaluation*

192 A survey was conducted in the study site inquiring about potato and fava bean crops (four
193 farmers of each crop type), including sowing and harvesting dates, management practices
194 and yields. Table 3 includes the characteristics of potato and fava bean plots, to which data
195 on sowing and harvesting dates, management practices and yields correspond.

196

197

198 Weather data (daily data on precipitation, maximum air temperature, minimum air
199 temperature and incoming solar radiation from 1997 to 2017) were collected from a station
200 located in the study area (Zahle).

201

202 **2.2.3. Model calibration and evaluation process**

203 Following Belhouchette et al. (2008), only the two most sensitive parameters were
204 calibrated for simulation with CropSyst, namely: the above ground biomass transpiration
205 coefficient (AGB_T) and the conversion of light to above ground biomass coefficient

206 (AGB_{IPAR}). These parameters were derived manually by changing the crop parameters until
 207 a satisfactory agreement between the predicted and simulated yield and biomass was
 208 achieved (Singh et al., 2008). While for potato and fava bean crops, the calibration was
 209 based on yields reported through the questionnaire conducted, as suggested by Komarek et
 210 al. (2017).

211 CropSyst was validated by comparing the simulated and measured values of the observed
 212 plots used for validation (section 2.2.2, Tables 1, 2 and 3), which were not part of the
 213 calibration process (one plot of each crop was used for calibration, while the others were
 214 used for validation). In the case of wheat, these values correspond to AGB, AGN, and SWC
 215 over the whole growing cycle, while for potato and fava bean crops, according Komarek
 216 et al. (2017), only the yield was validated after calibration. Root Mean Square Error
 217 (RMSE) was used to calculate the error of estimates as:

$$218 \quad RMSE = [Np^{-1} \sum_{i=1}^N (P_i - O_i)^2]^{0.5} \quad (3)$$

219 where Np is the number of pairs of observed (O_i) and simulated (P_i) data.

220 Then, the RMSE was computed relative to the mean of the observed values (\bar{O}) as:

$$221 \quad RRMSE = \frac{RMSE}{\bar{O}} \quad (4)$$

222 To have proper insight on the model efficiency, the model efficiency “EF” indicator was
 223 calculated as:

$$224 \quad EF = 1 - \frac{\sum_{i=1}^N (P_i - O_i)^2}{\sum_{i=1}^N (O_i - \bar{O})^2} \quad (5)$$

225 The model efficiency indicator varies from $-\infty$ to $+1$. Negative values can indicate bias in
226 linear models, yet could not be the case in non-linear models.

227 As an indicator to estimate correlation/regression, index of agreement “d” was calculated
228 as:

$$229 \quad d = 1 - \frac{\sum_{i=1}^N (P_i - O_i)^2}{\sum_{i=1}^N (|P_i - \bar{O}| + |O_i - \bar{O}|)^2} \quad (6)$$

230 where O_i represents the observed data, P_i represents the predicted data and \bar{O} is the average
231 of the observed data. The Willmott index of agreement (d) varies from 0.0 (poor model) to
232 1.0 (perfect model), similar to the interpretation of the coefficient of determination (R^2).

233 Eqs. 3, 4, 5 and 6 were applied to the validation plots within the region (four winter wheat
234 plots, three potato plots, and three fava bean plots), to make sure that by changing the
235 management practices and initial conditions, the model kept on generating satisfactory
236 estimates.

237

238 **2.3. Developing the scenarios to be simulated by the CropSyst model**

239 In this study, different wheat-based cropping systems of different rotation types (wheat-
240 wheat, wheat-fallow, wheat-potato, and wheat-fava bean, see Table 3) in two soil water
241 holding capacity (WHC) types, are simulated. The two soil types consist of different
242 horizons of different depth. Table 4 illustrates the soil characteristics of each soil type.
243 Each soil type is not uniform in terms of depth, rather consists of separate soil horizons.

244

245

246 We note hereby that full fertilization (Nitrogen) and full irrigation (water) follow the
247 amounts applied on the plot used for calibration (Plot ID 1 in Table 1), using the fixed
248 fertilization and fixed irrigation at fixed dates in CropSyst model. Four types of wheat
249 management systems were considered:

250 F1-I1: Full fertilization (230 kgN ha⁻¹) and full irrigation (100 mm)

251 F1-I0: Full fertilization (230 kgN ha⁻¹) and no irrigation.

252 F0-I1: No fertilization and full irrigation (100 mm).

253 F0-I0: No fertilization and no irrigation.

254 Thus, it is noted that during the simulations, these different management practices are only
255 applied in the case of wheat, while the management of potato and fava bean crops does not
256 change (management of Potato 1 and Fava bean 1 in Table 3 are used for simulation). As
257 shown by El Khansa (2017), potato farmers do not alter their water and fertilizer inputs as
258 they know in advance the high risk this involves. In the case of fava beans, the management
259 is fixed, a small amount of nitrogen is the only input they apply, if any.

260 Each cropping system scenario is run from 1997 to 2017. The output of each scenario
261 (cropping system) is 20 simulated years composed of 10 rotations (each lasts 2 years).
262 Table 5 summarizes the scenarios simulated. We note that the irrigation was simulated in
263 CropSyst using a fixed amount of water in a fixed time, as expressed by the farmers during
264 our survey.

265

266

267 **2.4. Calculation of the productivity and efficiency indicators for assessing the**
268 **performance of wheat-based cropping systems**

269 *Calculation of the productivity indicators*

270 - The net profit (US\$ ha⁻¹) indicator: computed at rotational level (2 years) as follows:

271
$$NP_r = Rev_r - Cost_r \quad (7)$$

272 Where *NP* stands for net profit at rotational level (*r*), *Rev* stands for revenue per rotation
273 (*r*) and *Cost* stands for the variable production cost of each rotation (*r*). We note hereby
274 that the variable production cost (*Cost*) represents all the costs needed to establish a
275 particular rotation. As the total simulation period is 20 years, the net profit is calculated 10
276 times for each cropping system.

277 In order to calculate the *Rev_r* and the *Cost_r* (Eq. 7) of each of the cropping systems, data on
278 input costs and output prices were collected through a local survey conducted at the study
279 site (Table 6), since in Lebanon, there is a lack of national official statistical sources for
280 annual input and output costs and prices. The input costs collected correspond to costs
281 related to cultivation (wheat, potato or fava bean), while output prices refer to selling the
282 produce (i.e. wheat grain yield, straw yield, potatoes, and fava bean grains). The input costs
283 and output prices collected through our survey correspond to an average of 5 years as the
284 prices are more or less stable and do not witness dramatic fluctuations. This also appears
285 when comparing to FAOSTAT (<http://www.fao.org/faostat/>) data. Regarding the
286 respondents, costs related to wheat were asked from 10 wheat farmers, costs related to
287 potato were asked from 10 potato farmers and costs related to fava bean were asked from
288 6 fava bean farmers. The farmers were selected randomly, yet we intended to interview

289 those who have been exercising the cultivation for a relatively long period (minimum 10
290 years). In addition, we aimed asking farmers who own their lands. The responses collected
291 from these farmers (Table 6) were homogeneous by looking at the averages and the
292 standard deviations.

293 It is noted that the input costs for both potato and fava bean crops are the variable
294 production costs including costs of fertilizer, water seeds, and labour. As simulating
295 different management systems is only carried out in the case of wheat, the variable
296 production cost (the only cost considered) to establish potato or fava bean cultivation is
297 equal to the input cost (Table 6) and does not change with the different scenarios.

298

299

300 - The average protein production (kg ha^{-1}) indicator is computed at rotational level
301 following two equations (Eqs. 8 & 9) for each cropping system. The purpose of this is to
302 serve comparing rotations of different crops within, to eventually come up with conclusions
303 on the protein production of each rotation.

304 First, the protein production (kg ha^{-1}) for each rotation is computed by considering the final
305 yield of each crop within the rotation and its corresponding protein percentage:

$$306 \quad P_r = [Yield_{Crop1} \times \%P_{Crop1}] + [Yield_{Crop2} \times \%P_{Crop2}] \quad (8)$$

307 where P_r is the amount of protein (kg ha^{-1}) produced by each rotation (r) (2years), $Yield_{Crop1}$
308 is the yield (kg ha^{-1}) of the first crop within the rotation, $Yield_{Crop2}$ is the yield (kg ha^{-1}) of
309 the second crop within the rotation, $\%P_{Crop1}$ is the percentage of protein contained in 1 kg

310 of the yield of crop1 and $\%P_{Crop2}$ is the percentage of protein contained in 1 kg of the yield
311 of crop2. Equation 8 is applied 10 times for each cropping system.

312 $Yield_{Crop1}$ and $Yield_{Crop2}$ are obtained from the CropSyst outputs. As for $\%P_{Crop1}$ and
313 $\%P_{Crop2}$, according to the USDA reports (USDA, 2018), 1 kg of grain contains 124.2 g of
314 protein, 1 kg of potato tuber_r contains 25.7 g of protein and 1 kg of fava bean_r contains
315 261.2 g of protein.

316 Then, the accumulated protein amounts of the 10 rotations (Nr) within each cropping
317 system are added up. Eventually, the average is measured by dividing the product by the
318 total number of rotations within each cropping system:

$$319 \quad \bar{P}_r = \frac{\sum_{r=1}^{Nr} (P_r)}{Nr} \quad (9)$$

320 where \bar{P}_r is the average protein production (kg ha^{-1}) at rotational level. r is the rotation. Nr
321 is the total number of rotations (=10 in this study).

322 ***Calculation of the efficiency indicators***

323 The nitrogen-use efficiency (NUE) (Darwish et al., 2006a; Gaudin et al., 2015;
324 Rahimizadeh et al., 2010) and water-use efficiency (WUE) (Kang et al., 2002; Sadras,
325 2004) for the average 10-year wheat crop presented in each of the cropping systems
326 (rotation type and wheat management) are computed following two equations (Eqs. 10 &
327 11).

$$328 \quad NUE_{Wheat} = \frac{AGN_1}{(N_{Supply1} + TSM)} \quad (10)$$

$$329 \quad WUE_{Wheat} = \frac{GY}{ET_{Actual}} \quad (11)$$

330 where NUE_{wheat} is the nitrogen-use efficiency calculated for the 10-years (out of a total of
 331 20 years) of the wheat crop. AGN_I (kgN ha⁻¹) stands for above ground nitrogen when wheat
 332 is fertilized. $N_{SupplyI}$ corresponds to the amount of supplied N fertilizer (kgN ha⁻¹), when
 333 applied. TSM corresponds to total soil mineralization rate accounting for N soil pool (kgN
 334 ha⁻¹). WUE_{wheat} is the water-use efficiency calculated for the 10years (out of a total of 20
 335 years) of wheat cultivation. GY corresponds to grain yield. ET_{Actual} stands for
 336 evapotranspiration (mm).

337 Then, the apparent recovery efficiency by difference ($ARED$) is compared following
 338 equations 12 & 13, to capture the added value of supplied N fertilizers and irrigation to
 339 wheat within the different cropping systems.

$$340 \quad ARED_N = \frac{AGN_1 - AGN_0}{N_{Supply1} - N_{Supply0}} \quad (12)$$

$$341 \quad ARED_W = \frac{GY_1 - GY_0}{W_{Supply1} - W_{Supply0}} \quad (13)$$

342 where $ARED_N$ and $ARED_W$ correspond to apparent recovery efficiency by difference, for
 343 nitrogen and water respectively. $N_{Supply0}$ corresponds to the amount of supplied N fertilizer
 344 (kgN ha⁻¹), when not supplied, which is equal to zero. AGN_0 (kgN ha⁻¹) corresponds to the
 345 above ground N when wheat is never fertilized. GY_I (kg ha⁻¹) and GY_0 (kg ha⁻¹) stand for
 346 grain yield under full irrigation and no irrigation at all, respectively. $W_{SupplyI}$ stands for the
 347 amount of water supplied as irrigation (mm). $W_{Supply0}$ is the amount of water supplied as
 348 irrigation (mm), when not supplied, which is equal to zero.

349 **2.5. Calculation of the “economic risk of low relative productivity”**

350 Taking into account that in the area (MENA region and many developing countries), access
351 to banks' credits or other credit institutions has hardly been established or has fallen in
352 disorder (Asseldonk et al., 2013), the risk calculation considered in this study is in line with
353 the farmers' concerns of being financially secured to keep on sustaining their cropping
354 system with no or low financial failure by mobilizing their net profit to invest in the rotation
355 that follows by covering its variable production cost. The financial failure considered here
356 is not being able to re-establish their rotations for preserving their livelihoods. In practice,
357 this is seen when the net profit of a particular rotation in year 1 is less than the variable cost
358 of the same rotation in year 2, meaning that the farmer who wishes to re-cultivate this
359 particular system, must mobilize external resources to increase the difference between the
360 net profit and the variable production cost. The risk calculation proposed is an original
361 procedure different from the standard calculations within the literature. The variation in the
362 risk in this study is basically related to yield variations as prices in the area are seen more
363 or less stable over the last years. In addition, it is important to mention that the risk
364 considered in this study is to compare different cropping systems (scenarios). In other
365 words, when the financial failure is seen, it does not mean that the farmer's livelihood is
366 terminated, yet they witness a risk of not being able independently in re-cultivating the
367 same system for the next rotation (two coming years), by covering its total production
368 costs.

369 Concretely, the economic risk of low relative productivity, which is expressed as a score,
370 is calculated as follows:

$$371 \text{Risk}_{CS} = (\sum_{r=1}^{Nr} W_r \times (+RD_r)) \times F_{r/Nr} \quad (14)$$

372 where $Risk_{cs}$ is the economic risk score that will be assigned to each of the cropping
373 systems, taking into account the 10 rotations within. r corresponds to the rotation. Nr
374 corresponds to the total number of rotations ($Nr=10$).

375 W_r is the normalized weight of the variable production cost at rotational level. As the tested
376 rotations have different production costs (different crops and management systems), a
377 normalized weight of the variable production cost is computed for each rotation, as a ratio
378 of the production costs of different rotations to the production cost of the most expensive
379 rotation (i.e. wheat-potato). Weight is calculated as follows:

$$380 \quad W_r = \frac{Cost_r}{Cost_{WP}} \quad (15)$$

381 where W_r is the weight of each rotation varying between 0 and 1, $Cost_r$ is the full production
382 cost of a rotation, $Cost_{WP}$ corresponds to the full production cost of the wheat-potato
383 rotation type.

384 RD_r is the relative deviation of the net profit from the cost at each rotation. Given that the
385 farmer will continue applying the same rotation type and wheat management system, the
386 relative deviation of this net profit/rotation from the total cost needed to re-establish the
387 same rotation (of the same rotation type and wheat management system) is computed as
388 follows:

$$389 \quad RD_r = \frac{Cost_r - NP_r}{Cost_r} \quad (16)$$

390 where RD_r is the relative deviation of the net profit from the cost for each rotation within
391 each cropping system, $Cost_r$ is the variable production cost of a particular rotation (e.g.
392 wheat-wheat in F1-I1 management system) and NP_r is the net profit of this particular

393 rotation within a particular cropping system. It is noted that within the whole period (20
394 years) of simulation in a particular cropping system (10 rotations), the $Cost_r$ is fixed and
395 doesn't change, while the NP_r changes for every rotation (2 years). If the output of RD_r is
396 negative, meaning that the NP_r is higher than the $Cost_r$, then the corresponding rotation is
397 neglected and is not considered when applying Eq. 14.

398 $F_{r/Nr}$ is the frequency of rotations (ratio from 0 to 1) whose net profit is lower than the
399 variable production cost, out of the 10 rotations (Nr). To highlight the repetition of rotations
400 with a deficit (positive RD_r), i.e. in which the net profit is lower than the cost, the ratio of
401 the occurrence of this "bad" event from the whole number of rotations ($Nr=10$) is computed
402 as follows:

$$403 \quad F_{r/Nr} = \frac{\#PositiveRD_r}{Nr} \quad (17)$$

404 where $\#PositiveRD_r$ is the number of rotations within a cropping system whose RD_r is
405 positive ($NP_r < Cost_r$). Eq. 17 is applied for each of the 16 cropping systems, in both low
406 and high WHC soils.

407

408 **3. Results**

409 **3.1. Calibration and validation of the CropSyst model**

410 The results of the validation of the CropSyst model are therefore generated after the
411 calibration. Following the rating proposed by Jamieson et al. (1991), the RRMSE ranged
412 between 9.2% and 12.7%, it can thus be considered as good to excellent simulation of dry
413 matter production (DMP). For above ground nitrogen (AGN) simulation, RRMSE ranged

414 from 7.7% to 25.0% and can be considered as average to good. In the case of the average
415 soil water content (SWC) simulation, RRMSE ranged from 21.0% to 34.3%. As regarding
416 the efficiency indicator (EF), for the DMP, the values ranged between 0.946 and 0.99,
417 between 0.9 and 0.99 for AGN and between -5.5 and -0.57 for SWC. Concerning the index
418 of agreement (the correlation/regression indicator), the values ranged between 0.990 and
419 0.996 for the DMP, between 0.96 and 0.997. As for the SWC, the index of agreement
420 produced values between 0.53 and 0.64. Hence, the calibrated model can be counted as
421 satisfactory in terms of simulating yield, water, and nitrogen cycles.

422 **3.2. Wheat grain yield as altered by the effects of rotation, management system,** 423 **and soil type**

424 The effects of different management systems, soil WHC, and rotations on wheat grain
425 yields (kg ha^{-1}) are compared for the different wheat-based systems, as shown in Table 7.
426 Before applying the mean separation test, we checked for homogeneity (using Chi-square
427 test) and normality (using Shapiro Wilk's W test) assumptions. For all our cases, the null
428 hypothesis was rejected by the Chi-square test confirming that the rotation types are linked
429 to the wheat grain yield (significant results) and non-significant for the Shapiro-Wilk's W
430 test, thus the normality assumption was checked. Tuckey test (2-way ANOVA analysis)
431 was used to its ability in reducing type 1 and 3 errors (Acutis et al., 2012)

432

433

434 In low WHC soil, wheat grain yields produced by a wheat-potato rotation were the highest
435 (with no clear effect of the rotation type). However, in all rotations, wheat grain yields

436 significantly decrease as the input amounts (fertilization and irrigation) decrease or are not
437 applied (e.g. in the case of a wheat-potato rotation it drops from 4515 kg ha⁻¹ in F1-I1 to
438 3433 kg ha⁻¹ in F0-I0).

439 When wheat is fertilized with no irrigation (F1-I0), wheat grain yields in a wheat-wheat
440 rotation (3787 kg ha⁻¹) are similar to those observed in a wheat-potato rotation (3757 kg
441 ha⁻¹), meaning that water stress in such a soil type (low WHC) is more significant than
442 nitrogen stress. This finding agrees with the results reported earlier (Huang et al., 2003).

443 When wheat is not fertilized (irrigated or not), wheat grain yields in wheat-fallow and
444 wheat-fava bean rotations are higher than those observed in a wheat-wheat rotation (e.g.
445 3873 kg ha⁻¹ in a wheat-fava bean rotation versus 3447 kg ha⁻¹ in a wheat-wheat rotation)
446 and similar to those observed in a wheat-potato rotation (3280 kg ha⁻¹ in a wheat-fava bean
447 rotation versus 3433 kg ha⁻¹ in a wheat-potato rotation). This means, indeed, that fallow
448 and fava bean when in rotation with wheat better mitigate water and nitrogen stresses on
449 wheat grain yields, in comparison with wheat-wheat rotation. This result is in agreement
450 with other findings (López-Bellido et al., 2012).

451 In high WHC soil, when wheat is fertilized (F1-I1 and F1-I0), there is no significant effect
452 of the rotation type on the wheat grain yield production. This is primarily attributed to the
453 type of soil (high WHC) that can hold more green water than low WHC. Hence, the loss in
454 yield that appeared in all the rotations from irrigated to rain-fed in high WHC (I1 to I0) is
455 due to water stress. Similar results were reported in the region (Sohi et al., 2009). However,
456 this loss is not as prominent as the one seen in low WHC soil (11% versus 18% of wheat
457 grain yield drop).

458 However, when nitrogen is limited (F0-I1 and F0-I0), as was reported for the soils of the
459 NENA region (Darwish et al., 2018), the wheat-wheat rotation always leads to significantly
460 lower wheat grain yields (4665 kg ha⁻¹) than the other tested rotation types (over 5200 kg
461 ha⁻¹). This is because wheat-fallow and wheat-fava bean rotations are less intensive in
462 terms of nitrogen demand, and do not neglect the nitrogen fixation ability of legumes
463 (Constantin et al., 2010; Moreau et al., 2012), as well as the fertilization of potatoes in the
464 wheat-potato rotation. This means that nitrogen and water stresses are higher in the case of
465 a wheat-wheat rotation when compared with other rotations.

466 Nevertheless, still in high WHC soils, wheat-wheat is the only rotation in which wheat
467 grain yields did drop significantly when changing from a wheat intensive management
468 system (F1-I1) (6083 kg ha⁻¹) to wheat semi-intensive management systems (F1-I0 and F0-
469 I1) (5250 kg ha⁻¹ and 5126 kg ha⁻¹, respectively), as well as when both nitrogen and water
470 were ceased (F0-I0) (4665 kg ha⁻¹). This suggests that avoiding wheat intensive
471 management systems (when in rotation with fallow, potato and fava bean) would not cause
472 water nor nitrogen stress, preserving the level of wheat grain yields.

473 **3.3. Nitrogen and water Apparent Recovery Efficiency by Difference (ARED)**

474 To grasp the added value of input resources (water and nitrogen) on wheat grain yields, the
475 Apparent Recovery Efficiency by Difference (ARED) for the two input resources (Keller
476 and Keller, 1995; Rao et al., 1992) is calculated, for wheat grain yields following different
477 previous crops (wheat, fallow, potato and fava bean) in the two soil water holding
478 capacities (low and high) [Eqs. (12) & (13) section 2.4.1]. ARED was computed for
479 nitrogen in both cases of irrigation (I1 and I0), as well as for irrigation in both cases of

480 nitrogen supply (F1 and F0) (Fig. 2). The separation of means was done using 2-way
481 ANOVA analysis (Tukey test).

482 Concerning fertilization, wheat grain yields increased the most because of nitrogen
483 fertilization (ARED_N) when cultivated in a wheat-wheat rotation, whether irrigated or not,
484 in low WHC soil (Fig. 2i & 2iii) (4.61 and 2.79, respectively) or in high WHC (Fig. 2ii &
485 2iv) (4.3 and 2.46, respectively). This means that for each 1 kg of added N, the increase in
486 grain yields for wheat in the wheat-wheat rotation is greater than that observed in other
487 rotations. In other words, wheat grain yields in a wheat-wheat rotation will be more
488 sensitive to lower fertilization, and thus more likely to decrease than wheat grain yields in
489 other rotations. The lowest slope observed for wheat grain yields was in the case of a wheat-
490 fava bean rotation, conforming to several studies that show that fava bean is an excellent
491 previous crop (Angus et al., 2015; Plaza-Bonilla et al., 2017; Yau et al., 2003), which partly
492 reduces the dependence of the main crop (winter wheat in this case) on nitrogen fertilization
493 (Voisin et al., 2013).

494 In the case of wheat-fallow and wheat-potato rotations, the results were more nuanced. In
495 low WHC soil, fallow land allows for a better use of N than potato as a previous crop, as
496 soil nutrients are rebalanced and soil biota is re-established (Plaza-Bonilla et al., 2017). On
497 the other hand, potato becomes a better previous crop than fallow land in high WHC soils.
498 This is due to the ability of these soils to store more water and facilitate the flux of nitrogen
499 to the roots by mass flow, for the next crop.

500 Concerning irrigation, the results are quite surprising as in low WHC (Fig. 2v), wheat in a
501 wheat-wheat rotation requires less water (in terms of irrigation) than in other rotations,
502 which contradicts other studies that show that wheat is more sensitive to irrigation (thus

503 more dependent) in a wheat-wheat rotation type (Gu et al., 2002; Musick et al., 1994;
504 Zhang and Oweis, 1999). This result can be explained by larger periods of fallow land than
505 in the case of wheat-wheat rotations (maximum of 3 months of fallow). Therefore leading
506 to larger amounts of evaporated water. This evaporated water, however, becomes less
507 important in high WHC (Fig. 2vi), in which more water is stored to its high capacity. In
508 coherence with the literature (Passioura and Angus, 2010), well-fertilized wheat (F1)
509 becomes more water reactive in a wheat-wheat rotation than other rotations. On the other
510 hand, wheat in a wheat-fava bean rotation becomes more reactive to water if nitrogen is
511 not applied. These results are consistent with multiple published studies, which state that
512 legumes are excellent previous crops, especially in a poorly fertilized system.

513 **3.4. Trends of the crops' yields (10 rotations) over the simulation period**

514 Out of the period of 20-years of simulation, 10 particular years (1998, 2000, 2002, 2004,
515 2006, 2008, 2010, 2012, 2014 and 2016) witnessed the cultivation of each of the three
516 crops (wheat, potato and fava beans). In Figure 3 below, the trends of each of the three
517 crops is represented, for both water holding capacity soils (WHC). In each year, four
518 agricultural practices were simulated (Table 5). Thus, as we aim to show the general trend,
519 the values of the four outputs of the four agricultural practices simulated were averaged.

520 The average wheat grain yield trend shows a slight increase until the 6th rotation (1998 till
521 2008), then a sharp drop until the 8th rotation (2012) then an increase afterwards. Similar
522 to the trend of winter wheat, the yield of fava beans increased from the 1st rotation till the
523 2nd one, then slight increase till the 6th rotation before the sharp drop till the 8th rotation
524 (2012). An increase was seen afterwards till the 9th rotation before finally a decrease at the
525 10th one. As for potato, the trend was more or less stable with a slight decrease until the 7th

526 rotation, continues decrease was seen till the 8th rotation before a sharp increase at the 9th
527 and 10th rotation.

528 **3.5. Rotation performance (productivity and efficiency) and economic risk of low** 529 **relative productivity**

530 Productivity (protein and net profit) at rotational level [Eqs. (7) & (9) section 2.4.1] versus
531 the resource-use efficiency calculated for wheat crops [Eqs. (10) & (11) section 2.4.1], in
532 each of the cropping systems are demonstrated (Fig. 4) for both soil types (low and high
533 WHC). In addition, based on the economic risk score calculated following Eq. 14 (section
534 2.4.2), the risk level of each of the cropping system is expressed.

535

536 Looking at the NUE (Fig. 4a & 3c), it is clear that for all systems efficiency decreases
537 dramatically when nitrogen is applied. With respect to protein production (Fig. 4a),
538 regardless of the management type, wheat-wheat (systems 1, 5, 9 and 13) and wheat-fava
539 bean (systems 4, 8, 12 and 16) rotations produce the highest amounts of protein (between
540 0.75 and 1.1 t rotation⁻¹ per ha followed by wheat-potato rotations (systems 3, 7, 11 and
541 15) that produce lower amounts depending on the management type (between 0.7 and 0.9
542 t rotation⁻¹ per ha). Wheat-fallow rotations (systems 2, 6, 10 and 14) produce the least
543 amount of protein (between 0.3 and 0.55 t rotation⁻¹ per ha). These results show that most
544 cropping systems, when grown in low WHC soil in such semi-arid areas, are over-
545 fertilized, which is relatively consistent with previous studies (Asseng et al., 2001; Ben
546 Zekri et al., 2018; Garabet et al., 1998). Residual soil nitrogen can be subject to nitrification
547 in well aerated loamy soils and nitrates can be leached and pollute the groundwater
548 (Darwish et al., 2003).

549 Regarding the net profit (Fig. 4c), it is obvious from the results that the wheat-potato
550 rotation is the most profitable rotation among the different rotations (from 8500
551 US\$/rotation/ha to 8700 US\$ rotation⁻¹ per ha). Wheat-fava bean (around 2000 US\$
552 rotation⁻¹ per ha) and wheat-wheat (around 1300 US\$ rotation⁻¹ per ha) rotations follow.
553 Eventually, wheat-fallow is the least profitable in terms of net profit (around 660 US\$
554 rotation⁻¹ per ha). This also confirms, therefore, that the cropping systems within the area
555 are over-fertilized since efficiency decreases while the net profit does not witness a similar
556 increase when intensifying the management system. In the long term, these practices may
557 affect soil-ecosystem functions.

558 The wheat-potato rotation is the riskiest one, compared to other rotations. In addition, the
559 wheat-fava bean rotation is not just more profitable than the wheat-wheat rotation, it is also
560 economically much less risky. Thus, growing legumes in rotation with wheat reduces
561 economic risk, as well as water and nitrogen dependence, compared to other rotations.
562 Similarly, and on a more general basis, in low WHC soil types, the more intensive the
563 systems, the riskier (economically) they are. This result contradicts several other studies
564 that suggest intensification, as a factor, to increase production stability (Gaudin et al., 2015;
565 Hartmann et al., 2015). This result shows that by intensifying the system in low WHC, that
566 is to say dry-land soils that are poor in terms of physical and biological properties,
567 productivity remains, efficiency decreases and economic and environmental sustainability
568 decrease.

569 By looking at the WUE, intensive systems (except wheat-fallow rotations) are more
570 efficient in terms of water-use than extensive systems. This result is consistent with the
571 literature, which mentions that water is one of the most limiting factors in shallow soils of

572 arid areas (Sultana et al., 2009). Semi-intensive and extensive systems (systems 8, 9, 11,
573 12, 13, 15 and 16) are then less efficient in terms of WUE. Wheat-fallow systems
574 (especially the extensive ones: 10 and 14) are the least efficient in terms of water, basically
575 due to large amounts of evaporated water.

576

577 By looking at the NUE (Fig. 5a & 5c), in high WHC soil, two groups of systems could be
578 observed, belonging to fertilized (low NUE) and unfertilized systems (high NUE). Within
579 the second group, wheat-fava bean (systems 4, 8, 12 and 16) and wheat-fallow (systems 2,
580 6, 10 and 14) rotations are superior to wheat-wheat (systems 1, 5, 9 and 13) and wheat-
581 potato (systems 3, 7, 11 and 15) rotations in terms of NUE. With respect to protein
582 production (Fig. 5a), wheat-wheat and wheat-fava bean rotations were the best rotations
583 compared to the other two rotations (1.2 to 1.5 t rotation⁻¹ per ha), followed by wheat-
584 potato (1.1 t rotation⁻¹ per ha) and wheat-fallow (0.7 to 0.8 t rotation⁻¹ per ha) rotations.
585 Results for soils with high WHC show that all rotations (except the wheat-wheat rotation)
586 are over-fertilized. As for the net profit, regardless of the soil type and WHC, the wheat-
587 potato rotation is the most profitable rotation (up to 12000 US\$ rotation⁻¹ per ha). Wheat-
588 fava bean and wheat-wheat rotations follow (3500 US\$/rotation/ha and 2500
589 US\$/rotation/ha, respectively). The wheat-fallow rotation comes last with around 1500
590 US\$/rotation/ha.

591 Comparing the results of high WHC to those of low WHC soil types, wheat-wheat and
592 wheat-potato rotations in high WHC soils become much less risky (if intensive
593 management in terms of nitrogen is avoided) in terms of economic risk of low productivity.

594 Wheat-fallow and wheat-fava bean rotations, similarly to low WHC soils, are the least
595 risky, economically, if adopted by farmers.

596 Regarding the WUE (Fig. 5b & 5d), the results show that water is not a limiting factor in
597 high WHC soils (Zhang et al., 2008). Even though fertilized systems have shown slightly
598 higher WUE than non-fertilized systems, the difference is not significant.

599

600 **4. Discussion**

601 When considering high resource-use efficiency, high system productivity (protein and
602 profit) as well as low economic risk in terms of system sustainability as a whole package,
603 it is clear that there is no comprehensible optimal scenario. Depending on our simulation
604 results, the productivity (protein and net profit) of the different wheat-based cropping
605 systems in two different soil WHC types, taking into consideration risk and wheat
606 efficiency results, are plotted on a conceptual guide-map (Fig. 6). In addition, the
607 productivity frontier is displayed to understand the best attainable scenarios.

608

609 Using this conceptual guide-map (Fig. 6) is essential for comparing the performance of the
610 different wheat-based cropping systems, but also to identify the possible levers to improve
611 the performance of these systems:

612 - Preserving deep soils for wheat cultivation: Such a strategy is achieved by combatting
613 ongoing soil degradation, especially in dry and sub-dry areas. The presence of many typical
614 cereal area plains in the Mediterranean region with low soil water holding capacity results
615 in grain yield reduction due to post-anthesis terminal drought where a strong relationship

616 was found between actual evapotranspiration in the grain filling phase and the final grain
617 yield. (Karrou and Oweis, 2012). As an example, the soil in the Medjerda-Tunisia (Souissi
618 et al., 2017), Saïs-Morocco (Mohamed et al., 2018) and the Bekaa plain of Lebanon
619 (Darwish et al., 2006b) typical cereal plains are more than 60% characterized by a low
620 water holding capacity.

621 Deep soils in arid area are currently mostly dominated by cereal crops (especially durum
622 and soft wheat), but a wide range of irrigated crops can also be observed, such as
623 vegetables, orchards and fodder crops which represent at least 30% of the total cultivated
624 area (Caiserman et al., 2019). These crops are less sensitive to the depth of the soil than
625 cereal crops. Therefore, keeping deep soils for cereals could be a leverage for policy-
626 makers in order to increase their production, input-use efficiency, and reduce the economic
627 risk of low relative productivity by at least 48%, 35% and 38% respectively, as shown in
628 this study.

629 - Reduction of the areas dedicated to wheat-fallow cropping systems (WF in Fig.5): such a
630 cropping system is characterized by low economic and nutritional performances compared
631 to other cropping systems. This explains why this system has gradually disappeared from
632 arid areas, and only exists for those practicing multiple activities (MoA, 2010). For those
633 farmers, the advantage of this system is that it requires very few inputs (particularly in high
634 WHC soils) and especially involves very little risk compared to other cropping systems.
635 Today, in the Mediterranean region, even if the areas dedicated to this type of rotation are
636 less common than those dedicated to other rotation types, half of the land is left
637 uncultivated each year (López-Bellido and López-Bellido, 2003). Most of the current
638 intensification policies in dry areas encourage the mobilization of more inputs by totally or

639 partially subsidizing access to irrigation water, fertilizers, seeds, etc. By promoting wheat-
640 based alternative systems other than wheat-fallow, it is potentially possible, according to
641 our results, to increase rotational (2 year) protein production by at least 50%, at the Bekaa
642 plain level.

643 - Intensification of wheat cultivation by increasing the amount of inputs: as expressed
644 before, this lever is the most used by policy-makers and farmers to increase wheat
645 production (Pala et al., 2007; Sadras, 2004). The guide-map (Fig. 6) shows that the
646 intensification by increasing inputs is not effective in all cropping systems and the effects
647 on rotational performance, efficiency and risk are not consistent with all cropping systems.
648 Unfortunately, most farmers manage wheat cultivation regardless of the rotation type
649 (Armengot et al., 2011), by considering intensified wheat management systems, presented
650 in a wheat-wheat rotation as a reference pathway to increase productivity (Balkovič et al.,
651 2014). Such means, which are encouraged by policy-makers, are not always reasonable as
652 the efficiency of wheat in utilizing the resources decreases dramatically in different dry
653 areas in the Mediterranean (Ben Zekri et al., 2018; Giménez et al., 2016; Ryan et al., 2007;
654 Souissi et al., 2017; Yau et al., 2003).

655 - Switching from wheat-wheat rotation to wheat-legume crops (fava bean in our study):
656 The wheat-legume rotation has shown better productivity than the wheat-wheat system,
657 significantly higher efficiency (nitrogen and water) and much lower economic risk. Such
658 findings were not very surprising as diversified rotations with catch crops usually yield
659 high NUE (Beaudoin et al., 2005; Hansen et al., 2015; Moreau et al., 2012). Growing winter
660 wheat with low inputs leads to a small sacrifice in terms of productivity, which may be a
661 reason why farmers prefer intensified wheat-wheat cropping systems. Apart from

662 fertilization, mechanization limitations in legumes cultivation and the dependence on
663 labour (weeding, harvesting) and establishments (e.g. storage warehouses), which would
664 be pricey if not already owned by the farmer, are also obstacles preventing farmers from
665 an easy switch to wheat-legume rotations. Moreover, fava beans production is a local
666 product, which is sold locally and thus linked to the national level. This could be attributed
667 to the low-trust that is given by farmers regarding national and local markets fearing
668 “unlawful speculation”. Instead, many of them would prefer investing in wheat cultivation
669 knowing in advance that the prices will not witness any change (even if negligible), as the
670 government buys the grain yield.

671 Indeed, wheat farmers in Lebanon as well as in the MENA region do over-fertilize their
672 crops for several reasons. First, farmers tend to believe that applying high rates of N would
673 decrease their economic risk by increasing the grain weight and the final grain yield, and
674 second because they do benefit from stable prices due to the support system at the national
675 level. This issue (over-fertilization) has been widely reported in Tunisia (Cheikh M’hamed
676 et al., 2014; Thabet et al., 2010), Egypt and Morocco (FAO, 2018). In another study, it was
677 also reported that the over nitrogen fertilization all over the Mediterranean is highly
678 impacting, negatively, the soil and water quality in the area in addition to reducing the
679 economic yield (the huge N input represents 15% of the input cost in Morocco) (Heng et
680 al., 2007). On the other hand, a study in Italy has suggested the increase of nitrogen
681 application to up to 200 kg ha⁻¹ (similar to the rates applied in the Bekaa plain) for higher
682 grain weight and better yield, taking into account the high environmental impacts (e.g.
683 nitrogen leaching) (Abad et al., 2004).

684 **Conclusions**

685 Broadly, as frequently cited, increased production as well as increasing the efficiency in
686 using the resources are the main requirements for feeding a vastly growing and changing
687 world (Godfray et al., 2010). Many farmers, who work under small profit margins seek
688 high production as a primary goal like many industrialized systems, which definitely trade
689 off against ecosystem values and environmental aspects (Foley et al., 2005). Such an
690 approach eventually leads to what we witness today in terms of negative environmental
691 consequences and resource depletion (Pimentel, 2005), as well as negative social impacts
692 (Marks et al., 2010). The results of our research showed that careful considerations should
693 be coupled with recommending a specific cropping system, especially at field level. No
694 optimal scenario was found (rotation and management) that may simultaneously guarantee
695 low risk, significant protein production, large net profit, and high resource-use efficiency
696 (NUE and WUE), at least for the rotations simulated in this study. Several studies have
697 analyzed one crop (or one cropping system) in relation to its productivity and efficiency.
698 Our results, nevertheless, by allowing the farmers and policy-makers to categorize existing
699 systems in terms of their performance and risk, indicate that at field level, a wheat-legume
700 rotation in which wheat is cultivated under semi-intensive and/or extensive agricultural
701 management is very recommended, especially for those who cannot bear high-risk systems,
702 securing both high efficiency in terms of resource-use and great protein production. In
703 future research, anticipating our results, we intend to upgrade this study to the farm level,
704 where more criteria and parameters may come across to propose a whole integrated system
705 that is profitable, non-risky and sustainable, overpowering food security deterioration and
706 nevertheless efficient in terms of resource-use efficiency.

707

708 **Acknowledgments**

709 The authors acknowledge the Conseil National de la Recherche Scientifique (CNRS-
710 Liban) for financially supporting the project, which was implemented in collaboration with
711 the Mediterranean Agronomic Institute of Montpellier, France (CIHEAM-IAM) and
712 IRSTEA (Montpellier, France). They declare no conflict of interest.

713 **References**

- 714 Abad, A., Lloveras, J., Michelena, A., 2004. Nitrogen fertilization and foliar urea effects
715 on durum wheat yield and quality and on residual soil nitrate in irrigated
716 Mediterranean conditions. *F. Crop. Res.* 87, 257–269.
717 <https://doi.org/10.1016/j.fcr.2003.11.007>
- 718 Acutis, M., Scaglia, B., Confalonieri, R., 2012. Perfunctory analysis of variance in
719 agronomy, and its consequences in experimental results interpretation. *Eur. J.*
720 *Agron.* 43, 129–135. <https://doi.org/10.1016/j.eja.2012.06.006>
- 721 Ahmad, A., Liu, M., Zhang, Y., Pei, H., Umair, M., Qi, Y., Shen, Y., 2017. Evaluation of
722 the CropSyst Model during Wheat-Maize Rotations on the North China Plain for
723 Identifying Soil Evaporation Losses. *Front. Plant Sci.* 8, 1–14.
724 <https://doi.org/10.3389/fpls.2017.01667>
- 725 Ahmed, G., Hamrick, D., Gereffi, G., 2014. Shifting Governance Structures in the Wheat
726 Value Chain : Implications for Food Security in the Middle East and North Africa.
727 *Centrorossidoria.It* 1–28.
- 728 Angus, J.F., Kirkegaard, J.A., Hunt, J.R., Ryan, M.H., Ohlander, L., Peoples, M.B., 2015.

729 Break crops and rotations for wheat. *Crop Pasture Sci.* 66, 523–552.
730 <https://doi.org/10.1071/CP14252>

731 Armengot, L., José-María, L., Blanco-Moreno, J.M., Bassa, M., Chamorro, L., Sans,
732 F.X., 2011. A novel index of land use intensity for organic and conventional farming
733 of Mediterranean cereal fields. *Agron. Sustain. Dev.* 31, 699–707.
734 <https://doi.org/10.1007/s13593-011-0042-0>

735 Asseldonk, M. Van, Burger, K., Maitre, E., Muller, B., 2013. Linking crop insurance and
736 rural credit.

737 Asseng, S., Turner, N.C., Keating, B.A., 2001. Analysis of water- and nitrogen-use
738 efficiency of wheat in a Mediterranean climate. *Plant Soil* 233, 127–143.

739 Balkovič, J., van der Velde, M., Skalský, R., Xiong, W., Folberth, C., Khabarov, N.,
740 Smirnov, A., Mueller, N.D., Obersteiner, M., 2014. Global wheat production
741 potentials and management flexibility under the representative concentration
742 pathways. *Glob. Planet. Change* 122, 107–121.
743 <https://doi.org/10.1016/j.gloplacha.2014.08.010>

744 Beaudoin, N., Saad, J.K., Van Laethem, C., Machet, J.M., Maucorps, J., Mary, B., 2005.
745 Nitrate leaching in intensive agriculture in Northern France: Effect of farming
746 practices, soils and crop rotations. *Agric. Ecosyst. Environ.* 111, 292–310.
747 <https://doi.org/10.1016/j.agee.2005.06.006>

748 Belhouchette, H., Braudeau, E., Hachicha, M., Donatelli, M., Mohtar, R.H., Wery, J.,
749 2008. Integrating spatial soil organization data with regional agricultural
750 management simulation model: a case study in northern Tunisia. *Am. Soc. Agric.*

751 Biol. Eng. 51, 1099–1109.

752 Ben Zekri, Y., Barkaoui, K., Marrou, H., Mekki, I., Belhouchette, H., Wery, J., 2018. On
753 farm analysis of the effect of the preceding crop on N uptake and grain yield of
754 durum wheat (*Triticum durum* Desf.) in Mediterranean conditions. Arch. Agron.
755 Soil Sci. 0, 1–16. <https://doi.org/10.1080/03650340.2018.1514111>

756 Bennett, D., Reynolds, M., Mullan, D., Izanloo, A., Kuchel, H., Langridge, P.,
757 Schnurbusch, T., 2012. Detection of two major grain yield QTL in bread wheat
758 (*Triticum aestivum* L.) under heat, drought and high yield potential environments.
759 Theor. Appl. Genet. 125, 1473–1485. <https://doi.org/10.1007/s00122-012-1927-2>

760 Berntsen, J., Olesen, J.E., Petersen, B.M., Hansen, E.M., 2006. Long-term fate of
761 nitrogen uptake in catch crops. Eur. J. Agron. 25, 383–390.
762 <https://doi.org/10.1016/j.eja.2006.07.006>

763 Braudel, F., 1990. La Méditerranée et le monde méditerranéen à l'époque de Philippe le
764 Bel Tome 1.

765 Brooks, E.S., Pan, W.L., Lamb, B.K., Stöckle, C.O., Maureira, F., Chi, J., O'Keeffe, P.T.,
766 Waldo, S., Pressley, S.N., Huggins, D.R., 2017. Carbon and Water Budgets in
767 Multiple Wheat-Based Cropping Systems in the Inland Pacific Northwest US:
768 Comparison of CropSyst Simulations with Eddy Covariance Measurements. Front.
769 Ecol. Evol. 5, 1–18. <https://doi.org/10.3389/fevo.2017.00050>

770 Caiserman, A., Dumas, D., Bennafla, K., Faour, G., Amiraslani, F., 2019. Application of
771 Remotely Sensed Imagery and Socioeconomic Surveys to Map Crop Choices in the
772 Bekaa Valley (Lebanon). Agriculture 9, 57.

773 <https://doi.org/10.3390/agriculture9030057>

774 Catchpole, W.R., Wheeler, C.J., 1992. Estimating plant biomass: A review of techniques.
775 *Aust. J. Ecol.* 17, 121–131. <https://doi.org/10.1111/j.1442-9993.1992.tb00790.x>

776 Cheikh M'hamed, H., Rezig, M., Naceur, M. Ben, 2014. Deficit Irrigation of Durum
777 Wheat (*Triticum durum* Desf): Effects on Total Dry Matter Production, Light
778 Interception and Radiation Use Efficiency Under Different Nitrogen Rates. *Sustain.*
779 *Agric. Res.* 4, 26. <https://doi.org/10.5539/sar.v4n1p26>

780 Constantin, J., Beaudoin, N., Laurent, F., Cohan, J.P., Duyme, F., Mary, B., 2011.
781 Cumulative effects of catch crops on nitrogen uptake, leaching and net
782 mineralization. *Plant Soil* 341, 137–154. <https://doi.org/10.1007/s11104-010-0630-9>

783 Constantin, J., Mary, B., Laurent, F., Aubrion, G., Fontaine, A., Kerveillant, P.,
784 Beaudoin, N., 2010. Effects of catch crops, no till and reduced nitrogen fertilization
785 on nitrogen leaching and balance in three long-term experiments. *Agric. Ecosyst.*
786 *Environ.* 135, 268–278. <https://doi.org/10.1016/j.agee.2009.10.005>

787 Dalal, R.C., Strong, W.M., Weston, E.J., Cooper, J.E., Wildermuth, G.B., Lehane, K.J.,
788 King, A.J., C. J. Holmes, 2001. Sustaining productivity of a Vertisol at Warra,
789 Queensland, with fertilisers, no-tillage, or legumes. *Aust. J. Exp. Agric.* 41.

790 Darwish, T., Atallah, T., Fadel, A., 2018. Challenges of soil carbon sequestration in the
791 NENA region. *Soil* 4, 225–235. <https://doi.org/10.5194/soil-4-225-2018>

792 Darwish, T., Atallah, T., Hajhasan, S., Chranek, A., 2003. Management of nitrogen by
793 fertigation of potato in Lebanon. *Nutr. Cycl. Agroecosystems* 67, 1–11.

794 <https://doi.org/10.1023/A:1025107202143>

795 Darwish, T.M., Atallah, T.W., Hajhasan, S., Haidar, A., 2006a. Nitrogen and water use
796 efficiency of fertigated processing potato. *Agric. Water Manag.* 85, 95–104.
797 <https://doi.org/10.1016/j.agwat.2006.03.012>

798 Darwish, T.M., Khawlie, M., Jomaa, I., Abou Daher, M., Awad, M., Masri, T., Shaban,
799 A., Faour, G., Bou Kheir, R., Abdallah, C., Haddad, T., 2006b. Soil map of
800 Lebanon: 1: 50 000. CNRS-Lebanon, Monogr. Ser. 4, 367.

801 Di Falco, S., Perrings, C., 2005. Crop biodiversity, risk management and the implications
802 of agricultural assistance. *Ecol. Econ.* 55, 459–466.
803 <https://doi.org/10.1016/j.ecolecon.2004.12.005>

804 El Khansa, M., 2017. Finding pathways for enhancing irrigated farming systems in
805 Lebanon. Université Montpellier.

806 FAO, 2018. World fertilizer trends and outlook to, World fertilizer trends.

807 Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin,
808 F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard,
809 E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N.,
810 Snyder, P.K., 2005. Global Consequences of Land Use. *Science* (80-.). 309, 570–
811 574.

812 Garabet, S., Wood, M., Ryan, J., 1998. Nitrogen and water effects on wheat yield in a
813 Mediterranean-type climate. I. Growth, water-use and nitrogen accumulation. *F.*
814 *Crop. Res.* 57, 309–318. [https://doi.org/10.1016/S0378-4290\(98\)00075-6](https://doi.org/10.1016/S0378-4290(98)00075-6)

815 Gaudin, A.C.M., Janovicek, K., Deen, B., Hooker, D.C., 2015. Wheat improves nitrogen
816 use efficiency of maize and soybean-based cropping systems. *Agric. Ecosyst.*
817 *Environ.* 210, 1–10. <https://doi.org/10.1016/j.agee.2015.04.034>

818 Giménez, C., Stöckle, C.O., Suárez-Rey, E.M., Gallardo, M., 2016. Crop yields and N
819 losses tradeoffs in a garlic-wheat rotation in southern Spain. *Eur. J. Agron.* 73, 160–
820 169. <https://doi.org/10.1016/j.eja.2015.11.016>

821 Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F.,
822 Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C., 2010. Food security: The
823 challenge of feeding 9 billion people. *Science* (80-.). 327, 812–818.
824 <https://doi.org/10.1126/science.1185383>

825 Gu, B., Kang, S., Liang, Y., Cai, H., Hu, X., Zhang, L., 2002. Effects of limited irrigation
826 on yield and water use efficiency of winter wheat in the Loess Plateau of China.
827 *Agric. Water Manag.* 55, 203–216. [https://doi.org/10.1016/s0378-3774\(01\)00180-9](https://doi.org/10.1016/s0378-3774(01)00180-9)

828 Hansen, E.M., Munkholm, L.J., Melander, B., Olesen, J.E., 2010. Can non-inversion
829 tillage and straw retention reduce N leaching in cereal-based crop rotations? *Soil*
830 *Tillage Res.* 109, 1–8. <https://doi.org/10.1016/j.still.2010.04.001>

831 Hansen, E.M., Munkholm, L.J., Olesen, J.E., Melander, B., 2015. Nitrate Leaching,
832 Yields and Carbon Sequestration after Noninversion Tillage, Catch Crops, and
833 Straw Retention. *J. Environ. Qual.* 44, 868. <https://doi.org/10.2134/jeq2014.11.0482>

834 Hartmann, T.E., Yue, S., Schulz, R., He, X., Chen, X., Zhang, F., Müller, T., 2015. Yield
835 and N use efficiency of a maize-wheat cropping system as affected by different
836 fertilizer management strategies in a farmer's field of the North China Plain. *F.*

837 Crop. Res. 174, 30–39. <https://doi.org/10.1016/j.fcr.2015.01.006>

838 Heng, L.K., Asseng, S., Mejahed, K., Rusan, M., 2007. Optimizing wheat productivity in
839 two rain-fed environments of the West Asia-North Africa region using a simulation
840 model. *Eur. J. Agron.* 26, 121–129. <https://doi.org/10.1016/j.eja.2006.09.001>

841 Huang, M., Dang, T., Gallichand, J., Goulet, M., 2003. Effect of increased fertilizer
842 applications to wheat crop on soil-water depletion in the Loess Plateau, China.
843 *Agric. Water Manag.* 58, 267–278. [https://doi.org/10.1016/S0378-3774\(02\)00086-0](https://doi.org/10.1016/S0378-3774(02)00086-0)

844 Jamieson, P.D., Porter, J.R., Wilson, D.R., 1991. A test of the computer simulation model
845 ARCWHEAT1 on wheat crops grown in New Zealand. *F. Crop. Res.* 27, 337–350.
846 [https://doi.org/10.1016/0378-4290\(91\)90040-3](https://doi.org/10.1016/0378-4290(91)90040-3)

847 Kang, S.Z., Zhang, L., Liang, Y.L., Hu, X.T., Cai, H.J., Gu, B.J., 2002. Effects of limited
848 irrigation on yield and water use efficiency of winter wheat in the Loess Plateau of
849 China. *Agric. Water Manag.* 55, 203–216. [https://doi.org/10.1016/s0378-](https://doi.org/10.1016/s0378-3774(01)00180-9)
850 [3774\(01\)00180-9](https://doi.org/10.1016/s0378-3774(01)00180-9)

851 Karrou, M., Oweis, T., 2012. Water and land productivities of wheat and food legumes
852 with deficit supplemental irrigation in a Mediterranean environment. *Agric. Water*
853 *Manag.* 107, 94–103. <https://doi.org/10.1016/j.agwat.2012.01.014>

854 Kehoe, D.P., 1988. Economics of agriculture on Roman imperial estates in North Africa.
855 Vandenhoeck Ruprecht, Göttingen Popul.

856 Keller, A.A., Keller, J., 1995. Effective Efficiency: A Water Use Efficiency Concept for
857 Allocating Freshwater Resources. 22. *Winrock Int.* 20.

858 Komarek, A.M., Bell, L.W., Whish, J.P.M., Robertson, M.J., Bellotti, W.D., 2015.
859 Whole-farm economic, risk and resource-use trade-offs associated with integrating
860 forages into crop-livestock systems in western China. *Agric. Syst.* 133, 63–72.
861 <https://doi.org/10.1016/j.agsy.2014.10.008>

862 Komarek, A.M., Drogue, S., Chenoune, R., Hawkins, J., Msangi, S., Belhouchette, H.,
863 Flichman, G., 2017. Agricultural household effects of fertilizer price changes for
864 smallholder farmers in central Malawi. *Agric. Syst.* 154, 168–178.
865 <https://doi.org/10.1016/j.agsy.2017.03.016>

866 López-Bellido, L., Muñoz-Romero, V., Benítez-Vega, J., Fernández-García, P., Redondo,
867 R., López-Bellido, R.J., 2012. Wheat response to nitrogen splitting applied to a
868 Vertisols in different tillage systems and cropping rotations under typical
869 Mediterranean climatic conditions. *Eur. J. Agron.* 43, 24–32.
870 <https://doi.org/10.1016/j.eja.2012.05.002>

871 López-Bellido, R.J., López-Bellido, L., 2003. Effects of crop rotation and nitrogen
872 fertilization on soil nitrate and wheat yield under rainfed Mediterranean conditions.
873 *Agronomie* 21, 509–516. <https://doi.org/10.1051/agro:2001140>

874 Macdonald, A.J., Poulton, P.R., Howe, M.T., Goulding, K.W.T., Powlson, D.S., 2005.
875 The use of cover crops in cereal-based cropping systems to control nitrate leaching
876 in SE England. *Plant Soil* 273, 355–373. <https://doi.org/10.1007/s11104-005-0193-3>

877 Mahmood, F., Belhouchette, H., Nasim, W., Shahzad, T., Hussain, S., Therond, O.,
878 Fahad, S., Refat Sultana, S., Wery, J., 2017. Economic and environmental impacts
879 of introducing grain legumes in farming systems of Midi-Pyrenees region (France):

880 A simulation approach. *Int. J. Plant Prod.* 11, 65–88.

881 Marks, A.R., Harley, K., Bradman, A., Kogut, K., Barr, D.B., Johnson, C., Calderon, N.,
882 Eskenazi, B., 2010. Organophosphate pesticide exposure and attention in young
883 Mexican-American children: The CHAMACOS study. *Environ. Health Perspect.*
884 118, 1768–1774. <https://doi.org/10.1289/ehp.1002056>

885 MoA, 2010. Resultats globaux du module de base du recensement de l’agriculture.

886 Mohamed, B., Mounia, K., Aziz, A., Ahmed, H., Rachid, B., Lotfi, A., 2018. Sewage
887 sludge used as organic manure in Moroccan sunflower culture: Effects on certain
888 soil properties, growth and yield components. *Sci. Total Environ.* 627, 681–688.
889 <https://doi.org/10.1016/j.scitotenv.2018.01.258>

890 Monzon, J.P., Sadras, V.O., Andrade, F.H., 2012. Modelled yield and water use
891 efficiency of maize in response to crop management and Southern Oscillation Index
892 in a soil-climate transect in Argentina. *F. Crop. Res.* 130, 8–18.
893 <https://doi.org/10.1016/j.fcr.2012.02.001>

894 Moreau, P., Ruiz, L., Raimbault, T., Vertès, F., Cordier, M.O., Gascuel-Odoux, C.,
895 Masson, V., Salmon-Monviola, J., Durand, P., 2012. Modeling the potential benefits
896 of catch-crop introduction in fodder crop rotations in a Western Europe landscape.
897 *Sci. Total Environ.* 437, 276–284. <https://doi.org/10.1016/j.scitotenv.2012.07.091>

898 Mueller, N.D., Gerber, J.S., Johnston, M., Ray, D.K., Ramankutty, N., Foley, J.A., 2012.
899 Closing yield gaps through nutrient and water management. *Nature* 490, 254–257.
900 <https://doi.org/10.1038/nature11420>

901 Musick, J.T., Jones, O.R., Stewart, B.A., Dusek, D.A., 1994. Water-yield relationships
902 for irrigated and dryland wheat in the U.S. southern plains. *Agron. J.*
903 <https://doi.org/10.2134/agronj1994.00021962008600060010x>

904 Nasrallah, A., Baghdadi, N., Mhawej, M., Faour, G., Darwish, T., Belhouchette, H.,
905 Darwich, S., 2018. A Novel Approach for Mapping Wheat Areas Using High
906 Resolution Sentinel-2 Images. *Sensors* 18, 2089. <https://doi.org/10.3390/s18072089>

907 Pala, M., Ryan, J., Zhang, H., Singh, M., Harris, H., 2007. Water-use efficiency of
908 wheat-based rotation systems in a Mediterranean environment. *Agric. Water Manag.*
909 93, 136–144. <https://doi.org/10.1016/j.agwat.2007.07.001>

910 Palosuo, T., Kersebaum, K.C., Angulo, C., Hlavinka, P., Moriondo, M., Olesen, J.E.,
911 Patil, R.H., Ruget, F., Rumbaur, C., Takáč, J., Trnka, M., Bindi, M., Çaldağ, B.,
912 Ewert, F., Ferrise, R., Mirschel, W., Şaylan, L., Šiška, B., Rötter, R., 2011.
913 Simulation of winter wheat yield and its variability in different climates of Europe:
914 A comparison of eight crop growth models. *Eur. J. Agron.* 35, 103–114.
915 <https://doi.org/10.1016/j.eja.2011.05.001>

916 Passioura, J.B., Angus, J.F., 2010. *Improving Productivity of Crops in Water-Limited*
917 *Environments*, 1st ed, *Advances in Agronomy*. Elsevier Inc.
918 [https://doi.org/10.1016/S0065-2113\(10\)06002-5](https://doi.org/10.1016/S0065-2113(10)06002-5)

919 Pimentel, D., 2005. ENVIRONMENTAL AND ECONOMIC COSTS OF THE
920 APPLICATION OF PESTICIDES PRIMARILY IN THE UNITED STATES.
921 *Environ. Dev. Sustain.* 7, 229–252. <https://doi.org/10.1007/s10668-005-7314-2>

922 Plaza-Bonilla, D., Nolot, J.M., Raffaillac, D., Justes, E., 2017. Innovative cropping

923 systems to reduce N inputs and maintain wheat yields by inserting grain legumes
924 and cover crops in southwestern France. *Eur. J. Agron.* 82, 331–341.
925 <https://doi.org/10.1016/j.eja.2016.05.010>

926 Rahimizadeh, M., Kashani, A., Zare-Feizabadi, A., Koocheki, A., Nassiri-Mahallati, M.,
927 2010. Nitrogen use efficiency of wheat as affected by preceding crop, application
928 rate of nitrogen and crop residues. *Aust. J. Crop Sci* 4, 363–368.

929 Rao, A.C.S., Smith, J.L., Parr, J.F., Papendick, R.I., 1992. Considerations in estimating
930 Nitrogen Recovery Efficiency by the difference and isotopic dilution methods.
931 *Fertil. Res.* 33, 209–217. <https://doi.org/10.1007/BF01050876>

932 Reynolds, S.G., 1970. The Gravimetric Method of Soil Moisture Determination. *J.*
933 *Hydrol.* 11, 258–273.

934 Rodriguez, I.R., Miller, G.L., 2000. Using a Chlorophyll Meter to Determine the
935 Chlorophyll Concentration, Nitrogen Concentration, and Visual Quality of St.
936 *Augustinegrass.* *HortScience* 35, 751–754.

937 Rötter, R.P., Palosuo, T., Kersebaum, K.C., Angulo, C., Bindi, M., Ewert, F., Ferrise, R.,
938 Hlavinka, P., Moriondo, M., Nendel, C., Olesen, J.E., Patil, R.H., Ruget, F., Takáč,
939 J., Trnka, M., 2012. Simulation of spring barley yield in different climatic zones of
940 Northern and Central Europe: A comparison of nine crop models. *F. Crop. Res.* 133,
941 23–36. <https://doi.org/10.1016/j.fcr.2012.03.016>

942 Ryan, J., Pala, M., Masri, S., Singh, M., Harris, H., 2007. Rainfed wheat-based rotations
943 under Mediterranean conditions: Crop sequences, nitrogen fertilization, and stubble
944 grazing in relation to grain and straw quality. *Eur. J. Agron.* 28, 112–118.

945 <https://doi.org/10.1016/j.eja.2007.05.008>

946 Sadras, V., 2002. Interaction between rainfall and nitrogen fertilisation of wheat in
947 environments prone to terminal drought: Economic and environmental risk analysis.
948 *F. Crop. Res.* 77, 201–215. [https://doi.org/10.1016/S0378-4290\(02\)00083-7](https://doi.org/10.1016/S0378-4290(02)00083-7)

949 Sadras, V.O., 2004. Yield and water-use efficiency of water- and nitrogen-stressed wheat
950 crops increase with degree of co-limitation. *Eur. J. Agron.* 21, 455–464.
951 <https://doi.org/10.1016/j.eja.2004.07.007>

952 Sieling, K., Kage, H., 2006. N balance as an indicator of N leaching in an oilseed rape -
953 winter wheat - winter barley rotation. *Agric. Ecosyst. Environ.* 115, 261–269.
954 <https://doi.org/10.1016/j.agee.2006.01.011>

955 Sieling, K., Stahl, C., Winkelmann, C., Christen, O., 2005. Growth and yield of winter
956 wheat in the first 3 years of a monoculture under varying N fertilization in NW
957 Germany. *Eur. J. Agron.* 22, 71–84. <https://doi.org/10.1016/j.eja.2003.12.004>

958 Sinclair, T.R., Rufty, T.W., 2012. Nitrogen and water resources commonly limit crop
959 yield increases, not necessarily plant genetics. *Glob. Food Sec.* 1, 94–98.
960 <https://doi.org/10.1016/j.gfs.2012.07.001>

961 Singh, A.K., Tripathy, R., Chopra, U.K., 2008. Evaluation of CERES-Wheat and
962 CropSyst models for water-nitrogen interactions in wheat crop. *Agric. Water
963 Manag.* 95, 776–786. <https://doi.org/10.1016/j.agwat.2008.02.006>

964 Sohi, S., Lopez-Capel, E., Krull, E., Bol, R., 2009. Biochar's roles in soil and climate
965 change : A review to guide future research. *CSIRO L. Water Sci. Rep.* 05/09.

966 Souissi, I., Boisson, J.M., Mekki, I., Therond, O., Flichman, G., Wery, J., Belhouchette,
967 H., 2017. Impact assessment of climate change on farming systems in the South
968 Mediterranean area: a Tunisian case study. *Reg. Environ. Chang.*
969 <https://doi.org/10.1007/s10113-017-1130-8>

970 Stockle, C.O., Donatelli, M., Nelson, R., 2003. CropSyst, a cropping systems simulation
971 model. *Eur. J. Agron.* 18, 289–307. [https://doi.org/Pii S1161-0301\(02\)00109-0](https://doi.org/Pii%20S1161-0301(02)00109-0)

972 Stockle, C.O., Martin, S.A., Campbell, G.S., 1994. CropSyst, a cropping systems
973 simulation model: Water/nitrogen budgets and crop yield. *Agric. Syst.* 46, 335–359.
974 [https://doi.org/10.1016/0308-521X\(94\)90006-2](https://doi.org/10.1016/0308-521X(94)90006-2)

975 Sultana, H., Ali, N., Iqbal, M.M., Khan, A.M., 2009. Vulnerability and adaptability of
976 wheat production in different climatic zones of Pakistan under climate change
977 scenarios. *Clim. Change* 94, 123–142. <https://doi.org/10.1007/s10584-009-9559-5>

978 Thabet, C., Chebil, A., Othmane, A., 2010. Improving nitrogen and water use efficiency
979 for wheat production in Mediterranean countries: Case of Tunisia. *New Medit* 9, 54–
980 58.

981 USDA, 2018. National Nutrient Database for Standard Reference.

982 Valle, P.S., Koesling *, M., Lien, G., Arntzen, H., Flaten, O., Ebbesvik, M., 2004. Risk
983 and risk management in organic and conventional cash crop farming in Norway.
984 *Food Econ. - Acta Agric. Scand. Sect. C* 1, 195–206.
985 <https://doi.org/10.1080/16507540410019692>

986 Voisin, A.-S., Guéguen, J., Huyghe, C., Jeuffroy, M.-H., Magrini, M.-B., Meynard, J.-M.,

987 Mougel, C., Pellerin, S., Pelzer, E., 2013. Legumes for feed, food, biomaterials and
988 bioenergy in Europe: a review. *Agron. Sustain. Dev.* 34, 361–380.
989 <https://doi.org/10.1007/s13593-013-0189-y>

990 Wright, B., Cafiero, C., 2010. Grain reserves and food security in MENA countries. UC
991 Berkeley Conf. Agric. Dev.

992 Yau, S.K., Bounejmate, M., Ryan, J., Baalbaki, R., Nassar, A., Maacaroun, R., 2003.
993 Barley-legumes rotations for semi-arid areas of Lebanon. *Eur. J. Agron.* 19, 599–
994 610. [https://doi.org/10.1016/S1161-0301\(03\)00006-6](https://doi.org/10.1016/S1161-0301(03)00006-6)

995 Zhang, H., Oweis, T., 1999. Water-yield relations and optimal irrigation scheduling of
996 wheat in the Mediterranean region. *Agric. Water Manag.* 38, 195–211.
997 [https://doi.org/10.1016/S0378-3774\(98\)00069-9](https://doi.org/10.1016/S0378-3774(98)00069-9)

998 Zhang, X., Chen, S., Sun, H., Pei, D., Wang, Y., 2008. Dry matter, harvest index, grain
999 yield and water use efficiency as affected by water supply in winter wheat. *Irrig. Sci.*
1000 27, 1–10. <https://doi.org/10.1007/s00271-008-0131-2>

1001





Fig. 1. Location of the Bekaa plain in Lebanon as well as the study area within the red tile








Figure 2. Apparent Recovery Efficiency by Difference (ARED) of nitrogen and irrigation of wheat crops, within each rotation, soil, and management type. a, b and c are symbols characterizing ARED which are significantly different or not regarding rotation.






Figure 3. General yield (kg ha^{-1}) trends of the three crops simulated over the 10 rotations in both WHC soils.




Figure 4. Risk representation of each of the cropping systems in low WHC soil denoted by the scale bar from dark blue (not risky at all) to dark red (very risky). The variation of protein production at rotational level versus wheat NUE (a) and wheat WUE (b), the variation of net profit at rotational level versus wheat NUE (c) and wheat WUE (d). A legend for each ID is presented to the right of the figure.




Figure 5. Risk representation of each of the cropping systems in high WHC soil denoted by the scale bar from dark blue (not risky at all) to dark red (very risky). The variation of protein production at rotational level versus wheat NUE (a) and wheat WUE (b), the variation of net profit at rotational level versus wheat NUE (c) and wheat WUE (d). A legend for each ID is presented to the right of the figure.




Fig. 6. Conceptual guide-map defining the behavior of the most widespread cropping systems taking into account the inputs of the wheat crop, rotational outputs, wheat efficiency, and economic risk of low relative productivity. WW, WF, WP and WFB correspond to wheat-wheat, wheat-fallow, wheat-potato, and wheat-fava bean rotations, respectively. The darker the represented system area, the more intensive the management system.

Table 1. Winter wheat reference plot characteristics.

Plot ID	Sowing date	Harvesting date	Sowing density (Seeds m ⁻²)	N applied (kg ha ⁻¹)	Irrigation applied (mm)	Soil WHC
1	23/11/2018	02/07/2018	480	230	100	low
2	20/11/2018	05/07/2018	480	170	240	high
3	20/11/2018	01/07/2018	480	170	160	low
4	15/11/2018	01/07/2018	430	280	90	high
5	23/11/2018	28/07/2018	465	180	0	high

Table 2. Mean and standard deviation results of the in-situ measurements of wheat plots. The measurements correspond to above ground biomass (AGB), above ground nitrogen (AGN), and soil water content (SWC). The measurements took place at 5 physiological stages (1: sowing, 2: tillering, 3: booting, 4: flowering and 5: physiological maturity).

Plot ID	Physiological Stage	AGB (kg ha ⁻¹)		AGN (kgN ha ⁻¹)		SWC (m ³ m ⁻³)	
		Mean	SD	Mean	SD	SD	Mean
1	1					0.05	0.39
1	2	390	138	28.2	2.8	0.02	0.39
1	3	2527	126			0.02	0.4
1	4	6410	537	196.7	8.2	0.08	0.38
1	5	9975	722			0.05	0.32
2	1					0.03	0.26
2	2	513	30.5	26.2	1.6	0.05	0.36
2	3	2513	533			0.02	0.35
2	4	6290	377	287	17	0.02	0.34
2	5	10852	577			0.03	0.33
3	1					0.07	0.26
3	2	600	56	28.8	2.7	0.05	0.35
3	3	2817	241			0.03	0.34
3	4	5987	163	250.2	6.8	0.03	0.36
3	5	11338	463			0.02	0.36
4	1					0.03	0.26
4	2	529	58	21.1	2.3	0.03	0.29
4	3	2330	261			0.02	0.34
4	4	5557	681	222.3	27.2	0.02	0.29
4	5	8050	349			0.03	0.26
5	1					0.03	0.41
5	2	660	65	23.8	2.3	0.02	0.4
5	3	2980	356			0.02	0.38
5	4	5343	218	224.1	9.1	0.03	0.3
5	5	8962	730			0.03	0.28

Table 3. Potato and fava bean plot characteristics. The yield is expressed in dry matter (at a standard level of moisture).

Crop and ID	Sowing date	Harvesting date	N applied (kg ha ⁻¹)	Irrigation applied (mm)	Yield (kg ha ⁻¹)
Potato 1	05/03/2018	10/07/2018	100	560	50000
Potato 2	05/03/2018	15/07/2018	193	600	37000
Potato 3	01/03/2018	21/06/2018	370	700	40000
Potato 4	10/03/2018	17/07/2018	370	500	40000
Fava bean 1	10/11/2017	05/05/2018	50	50	1201
Fava bean 2	08/11/2017	15/05/2018	50	0	945
Fava bean 3	12/11/2017	11/05/2018	0	0	1125

Table 4 Soil characteristics in where the pilot fields were selected in the mid Bekaa plain of Lebanon.

Low	Depth	Soil Class	Sand %	Clay %	OM%	BD (g	WP	FC
	(m)					cm ⁻³)	(%Vol)	(%Vol)
Horizon 1	0.1	Cambisols	30	56	2.3	1.29	33.2	44.6
Horizon 2	0.25	Cambisols	25	55	2.2	1.28	32.6	44.1
Horizon 3	0.3	Cambisols	18	56	1.8	1.35	32.8	44.2
Horizon 4	0.4	Cambisols	25	48	1.9	1.33	33	44.4
High								
Horizon 1	0.1	Cambisols	29	55	2.6	1.3	32.6	44.1
Horizon 2	0.35	Cambisols	25	55	2.8	1.32	32.5	44
Horizon 3	0.55	Cambisols	16	50	2.2	1.28	29.6	42.2
Horizon 4	0.4	Cambisols	25	45	2	1.34	27.1	40.1

Table 5. Cropping system scenarios simulated using CropSyst. Each cropping system scenario consists of a rotation type and wheat management system. 16 cropping systems were simulated in two soil water holding capacity types, leading to 32 scenarios.

Low Soil holding capacity (LSC)			High soil holding capacity (HSC)		
Scenario	Rotation	Management	Scenario	Rotation	Management
1	Wheat-Wheat	F1-I1	17	Wheat-Wheat	F1-I1
2	Wheat-Wheat	F1-I0	18	Wheat-Wheat	F1-I0
3	Wheat-Wheat	F0-I1	19	Wheat-Wheat	F0-I1
4	Wheat-Wheat	F0-I0	20	Wheat-Wheat	F0-I0
5	Wheat-Fallow	F1-I1	21	Wheat-Fallow	F1-I1
6	Wheat-Fallow	F1-I0	22	Wheat-Fallow	F1-I0
7	Wheat-Fallow	F0-I1	23	Wheat-Fallow	F0-I1
8	Wheat-Fallow	F0-I0	24	Wheat-Fallow	F0-I0
9	Wheat-Potato	F1-I1	25	Wheat-Potato	F1-I1
10	Wheat-Potato	F1-I0	26	Wheat-Potato	F1-I0
11	Wheat-Potato	F0-I1	27	Wheat-Potato	F0-I1
12	Wheat-Potato	F0-I0	28	Wheat-Potato	F0-I0
13	Wheat-Fava bean	F1-I1	29	Wheat-Fava bean	F1-I1
14	Wheat-Fava bean	F1-I0	30	Wheat-Fava bean	F1-I0
15	Wheat-Fava bean	F0-I1	31	Wheat-Fava bean	F0-I1

16	Wheat-Fava bean	F0-I0	32	Wheat-Fava bean	F0-I0
----	-----------------	-------	----	-----------------	-------

Table 6. Crop input costs and output prices.

Input	Cost (US\$ ha ⁻¹)	Output	Price (US\$ Ton ⁻¹)	Crop	Input cost (US\$ ha ⁻¹)	Output price (US\$ Ton ⁻¹)
Wheat						
Fertilizers	400 (SD=47)	Grain yield	360	Potato	9150 (SD= 1556)	330 (SD=43)
Water	450 (SD= 48)	Straw	50 (SD=8)	Fava bean	745 (SD= 78)	1000 (SD=114)
Labor+ Pesticides	450 (SD= 70)					
Seeds	200 (SD= 31)					

Table 7. Wheat grain yields in different soil water holding capacities (WHC), rotations and management systems in dry Mediterranean conditions. The statistically different groups are represented by different letters (a, b and c) characterizing yields with significant difference (Tukey test at $\alpha=0.05$).

Management	Wheat grain yield (kg ha ⁻¹)				ANOVA Significant difference (Rotation)
	Wheat-Wheat	Wheat-Fallow	Wheat-Potato	Wheat-Fava bean	
Soil with low WHC					
F1-I1	4513	4374	4515	4404	a,b,a,ab
F1-I0	3787	3517	3757	3601	a,b,a,b
F0-I1	3447	3705	3962	3873	a,ab,b,b
F0-I0	3124	3141	3433	3280	a,a,b,ab
<i>Significant difference (Management)</i>	a,b,bc,c	a,b,b,b	a,b,ab,b	a,bc,ab,c	
Soil with high WHC					
F1-I1	6083	6246	6216	6117	a,a,a,a
F1-I0	5250	5638	5549	5525	a,a,a,a
F0-I1	5126	5818	5763	5736	a,b,b,b
F0-I0	4665	5421	5290	5200	a,b,b,b
<i>Significant difference (Management)</i>	a,b,b,b	a,ab,ab,b	a,ab,ab,b	a,ab,ab,b	